

ASYMPTOTIC ENERGY OF LATTICES

WEIGEN YAN AND ZUHE ZHANG

ABSTRACT. The energy of a simple graph G arising in chemical physics, denoted by $\mathcal{E}(G)$, is defined as the sum of the absolute values of eigenvalues of G . As the dimer problem and spanning trees problem in statistical physics, in this paper we propose the energy per vertex problem for lattice systems. In general for a type of lattices in statistical physics, to compute the entropy constant with toroidal, cylindrical, Mobius-band, Klein-bottle, and free boundary conditions are different tasks with different hardness and may have different solution. We show that the energy per vertex of plane lattices is independent on the toroidal, cylindrical, Mobius-band, Klein-bottle, and free boundary conditions. Particularly, the asymptotic formulae of energies of the triangular, $3^3.4^2$, and hexagonal lattices with toroidal, cylindrical, Mobius-band, Klein-bottle, and free boundary conditions are obtained explicitly.

1. INTRODUCTION

Throughout this paper, we suppose that $G = (V(G), E(G))$ is a simple graph with the vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and the edge set $E(G)$, if not specified. The adjacency matrix of G with n vertices, denoted by $A(G) = (a_{ij})_{n \times n}$, is an $n \times n$ symmetric matrix such that $a_{ij} = 1$ if vertices v_i and v_j are adjacent and 0 otherwise. The characteristic polynomial of G , denoted by $\phi(G, x)$, is defined as $\det(xI_n - A(G))$, where I_n is an identity matrix of order n . Denote the degree of vertex v_i of G by $d_G(v_i)$. If H is a subgraph of G , then $G - E(H)$ denotes the subgraph obtained from G by deleting all edges in H .

Gutman [6, 7] defined the energy of a graph G with n vertices, denoted by $\mathcal{E}(G)$, as $\mathcal{E}(G) = \sum_{i=1}^n |\lambda_i(G)|$, where $\lambda_i(G)$ are the eigenvalues of the adjacency matrix of G .

In statistical physics the dimer problem considers the molecular freedom (free energy per dimer) (see for example [5, 9, 10, 11, 13, 15]), the spanning tree problem considers the entropy of spanning trees [2, 14, 17], and the independent set problem considers the entropy of independent sets [1]. It is natural to consider the chemical physics parameter -

Key words and phrases. Energy, lattice, tensor product, characteristic polynomial.

The first author was supported in part by NSFC Grant(10771086) and by Program for New Century Excellent Talents in Fujian Province University.

the energy of lattices from the statistical physics point of view. For the case of quadratic lattices, to compute the energy per vertex is an easy task as follows.

Suppose that G_n, G'_n , and G''_n are the plane square lattices with toroidal, cylindrical, and free boundary conditions, respectively. That is, $G_n = C_n \times C_n, G'_n = P_n \times C_n$, and $G''_n = P_n \times P_n$, where C_n and P_n denote the cycle and the path with n vertices, and $G \times H$ is the Cartesian product of two graphs G and H . Obviously, $\{G''_n\}$ is a sequence of spanning subgraphs of the sequence $\{G'_n\}$ of finite graphs, and $\{G'_n\}$ is a sequence of spanning subgraphs of the sequence $\{G_n\}$ of finite graphs. Particularly,

$$\lim_{n \rightarrow \infty} \frac{|\{v \in V(G''_n) : d_{G''_n}(v) = d_{G_n}(v)\}|}{|V(G_n)|} = \lim_{n \rightarrow \infty} \frac{|\{v \in V(G'_n) : d_{G'_n}(v) = d_{G_n}(v)\}|}{|V(G_n)|} = 1,$$

that is, almost all vertices of G_n and G''_n (resp. G_n and G'_n) have the same degrees. On the other hand, it is well known that the eigenvalues of G''_n (resp. G'_n and G_n) are $2 \cos \frac{i\pi}{n+1} + 2 \cos \frac{j\pi}{n+1}, i, j = 1, 2, \dots, n$ (resp. $2 \cos \frac{i\pi}{n+1} + 2 \cos \frac{2j\pi}{n}, i = 1, 2, \dots, n, j = 0, 1, 2, \dots, n-1$, and $2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{n}, i, j = 0, 1, \dots, n-1$). Hence the energy per vertex of G''_n, G'_n , and G_n are defined as

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G''_n)}{|V(G''_n)|} &= \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G''_n)}{n^2} = \lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i,j=1}^n (|2 \cos \frac{i\pi}{n+1} + 2 \cos \frac{j\pi}{n+1}|) \\ &= 2 \int_0^1 \int_0^1 |\cos \pi x + \cos \pi y| dx dy = \frac{2}{\pi^2} \int_0^\pi \int_0^\pi |\cos x + \cos y| dx dy \approx 1.6211; \\ \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G'_n)}{|V(G'_n)|} &= \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G'_n)}{n^2} = \lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i=1}^n \sum_{j=0}^{n-1} (|2 \cos \frac{i\pi}{n+1} + 2 \cos \frac{2j\pi}{n}|) \\ &= 2 \int_0^1 \int_0^1 |\cos \pi x + \cos 2\pi y| dx dy = \frac{1}{\pi^2} \int_0^\pi \int_0^{2\pi} |\cos x + \cos y| dx dy \approx 1.6211; \\ \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G_n)}{|V(G_n)|} &= \lim_{n \rightarrow \infty} \frac{\mathcal{E}(G_n)}{n^2} = \lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i,j=0}^{n-1} (|2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{n}|) \\ &= 2 \int_0^1 \int_0^1 |\cos 2\pi x + \cos 2\pi y| dx dy = \frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{2\pi} |\cos x + \cos y| dx dy \approx 1.6211, \end{aligned}$$

implying G_n, G'_n , and G''_n have the same asymptotic energy ($\approx 1.6211n^2$).

The phenomenon above is not accidental. In this paper we obtain the asymptotic formulae of energies of triangular, $3^3.4^2$, and hexagonal lattices with toroidal, cylindrical, Mobius-band, Klein-bottle, and free boundary conditions. Our approach implies that in general the energy per vertex of plane lattices is independent of the boundary conditions.

2. THE ASYMPTOTIC ENERGIES OF SOME LATTICES

2.1. Graph asymptotic energy change due to edge deletion. Let us record the following results firstly. Koolen and Moulton [12] proved that if $2m \geq n$ and G is a graph on n vertices with m edges, then

$$\mathcal{E}(G) \leq \frac{2m}{n} + \sqrt{(n-1) \left[2m - \left(\frac{2m}{n} \right)^2 \right]}, \quad (1)$$

and if $2m \leq n$ and G is a graph on n vertices with m edges, then

$$\mathcal{E}(G) \leq 2m. \quad (1')$$

The following result is immediate from (1) and (1').

Proposition 2.1. *Let G be a graph with m edges. Then*

$$\mathcal{E}(G) \leq 2m.$$

Day and So [3, 4] first studied how the energy of a graph changes when edges are removed. They found the following

Lemma 2.2 (Day and So [3, 4]). *Let H be an induced subgraph of a graph G . Then*

$$\mathcal{E}(G) - \mathcal{E}(H) \leq \mathcal{E}(G - E(H)) \leq \mathcal{E}(G) + \mathcal{E}(H).$$

With a similar method, we can prove the following:

Lemma 2.3. *Let H be a subgraph of a graph G . Then*

$$|\mathcal{E}(G) - \mathcal{E}(H)| \leq \mathcal{E}(G - E(H)) \leq \mathcal{E}(G) + \mathcal{E}(H).$$

Given two graphs G and H ($V(G) \cap V(H)$ may be disjoint), Denoted by $\Delta(G, H) = |E(G)| + |E(H)| - 2|E(G) \cap E(H)|$, i.e., $\Delta(G, H)$ equals the number of edges of symmetric difference of $E(G)$ and $E(H)$.

Theorem 2.4. *Suppose $\{G_n\}$ and $\{H_n\}$ are two sequences of graphs such that*

$$\lim_{n \rightarrow \infty} \frac{\Delta(G_n, H_n)}{\mathcal{E}(G_n)} = 0.$$

Then

$$\lim_{n \rightarrow \infty} \frac{\mathcal{E}(H_n)}{\mathcal{E}(G_n)} = 1.$$

Proof. Let F_n be the subgraph of G_n or H_n induced by $E(G_n) \cap E(H_n)$. Note that

$$\begin{aligned} \left| \frac{\mathcal{E}(H_n)}{\mathcal{E}(G_n)} - 1 \right| &= \left| \frac{\mathcal{E}(H_n) - \mathcal{E}(G_n)}{\mathcal{E}(G_n)} \right| = \left| \frac{\mathcal{E}(H_n) - \mathcal{E}(F_n) + \mathcal{E}(F_n) - \mathcal{E}(G_n)}{\mathcal{E}(G_n)} \right| \\ &\leq \left| \frac{\mathcal{E}(G_n) - \mathcal{E}(F_n)}{\mathcal{E}(G_n)} \right| + \left| \frac{\mathcal{E}(H_n) - \mathcal{E}(F_n)}{\mathcal{E}(G_n)} \right|. \end{aligned}$$

By Lemma 2.3 and Proposition 2.1,

$$|\mathcal{E}(G_n) - \mathcal{E}(F_n)| \leq \mathcal{E}(G_n - E(F_n)) \leq 2|E(G_n)| - 2|E(F_n)|,$$

$$|\mathcal{E}(H_n) - \mathcal{E}(F_n)| \leq \mathcal{E}(H_n - E(F_n)) \leq 2|E(H_n)| - 2|E(F_n)|.$$

Hence

$$\left| \frac{\mathcal{E}(H_n)}{\mathcal{E}(G_n)} - 1 \right| \leq \frac{2\Delta(G_n, H_n)}{\mathcal{E}(G_n)}$$

implying the theorem. $\square$

Corollary 2.5. *Suppose that $\{G_n\}$ is a sequence of finite simple graphs with bounded average degree such that $\lim_{n \rightarrow \infty} |V(G_n)| = \infty$ and $\lim_{n \rightarrow \infty} \frac{\mathcal{E}(G_n)}{|V(G_n)|} = h \neq 0$. If $\{G'_n\}$ is a sequence of spanning subgraphs of $\{G_n\}$ such that $\lim_{n \rightarrow \infty} \frac{|\{v \in V(G'_n) : d_{G'_n}(v) = d_{G_n}(v)\}|}{|V(G_n)|} = 1$, then $\lim_{n \rightarrow \infty} \frac{\mathcal{E}(G'_n)}{|V(G'_n)|} = h$. That is, G_n and G'_n have the same asymptotic energy.*

A direct sequence of Corollary 2.5 is that $P_n \times P_n$, $P_n \times C_n$, and $C_n \times C_n$ have the same asymptotic energy which is shown in the introduction. More generally, by Corollary 2.5, we have

Remark 2.6. Suppose $G_i = P_n$ or $G_i = C_n$, $i = 1, 2, \dots, k$, and k is a constant. If n is sufficiently large, then the asymptotic energy of the n -dimensional lattices

$$\mathcal{E}(G_1 \times G_2 \times \dots \times G_k) \approx \frac{2n^k}{\pi^k} \int_0^\pi \int_0^\pi \dots \int_0^\pi \left(\sum_{i=1}^k |\cos x_i| \right) dx_1 dx_2 \dots dx_k.$$

Remark 2.7. Corollary 2.5 gives a method to calculate the asymptotic energy of a graph with bounded average degree. Suppose that $\{G_n\}$ is a sequence of finite simple graphs with bounded average degree. It is difficult to calculate its asymptotic energy directly. We can find a graph G'_n with bounded average degree, which satisfies $|V(G_n)| = |V(G'_n)|$ and almost all vertices of G_n and G'_n have the same degrees. If we can compute the asymptotic energy of G'_n directly, then by Corollary 2.5, G_n and G'_n have the same asymptotic energy. We will use this idea to calculate the asymptotic energy of some graphs in the next subsections.




FIGURE 1. (a) The triangular lattice $T^t(n, m)$ with toroidal boundary condition; (b) the triangular lattice $T^c(n, m)$ with cylindrical boundary condition; (c) the triangular lattice $T^f(n, m)$ with free boundary condition.

2.2. The triangular lattice. The triangular lattice with toroidal boundary condition, denoted by $T^t(n, m)$, can be regarded as an $n \times m$ square lattice with toroidal boundary condition with an additional diagonal edge added, in the same way, to every square, see Figure 1(a), where $(a_1, a_1^*), (a_2, a_2^*), \dots, (a_m, a_m^*); (b_1, b_1^*), (b_2, b_2^*), \dots, (b_n, b_n^*); (b_1^*, b_2), (b_2^*, b_3), \dots, (b_{n-1}^*, b_n), (b_n^*, b_1) = (a_m^*, a_1); (a_1^*, a_2), (a_2^*, a_3), \dots, (a_{m-1}^*, a_m)$ are edges in $T^t(n, m)$. If we delete the edges $(b_1, b_1^*), (b_2, b_2^*), \dots, (b_n, b_n^*); (b_1^*, b_2), (b_2^*, b_3), \dots, (b_{n-1}^*, b_n), (b_n^*, b_1)$ from $T^t(n, m)$, the triangular lattice with cylindrical boundary condition, denoted by $T^c(n, m)$, is obtained (see Figure 1(b)). If we delete the edges $(a_1, a_1^*), (a_2, a_2^*), \dots, (a_m, a_m^*), (a_1^*, a_2), (a_2^*, a_3), \dots, (a_{m-1}^*, a_m)$ from $T^c(n, m)$, the triangular lattice with free boundary condition, denoted by $T^f(n, m)$ is obtained (see Figure 1(c)). The asymptotic number of perfect matchings of $T^t(n, m)$ can be found in Wu [15]. The asymptotic number of spanning trees of $T^t(n, m)$ (resp. $T^c(n, m)$ and $T^f(n, m)$) was obtained by Shrock and Wu [14] (resp. by Yan and Zhang [17]).

Theorem 2.8. *For the triangular lattices $T^t(n, m)$, $T^c(n, m)$, and $T^f(n, m)$ with toroidal, cylindrical, and free boundary conditions,*

$$\begin{aligned} \lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(T^t(n, m))}{nm} &= \lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(T^c(n, m))}{nm} = \lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(T^f(n, m))}{nm} \\ &= \frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{2\pi} |\cos x + \cos y + \cos(x + y)| dx dy (\approx 2.065), \end{aligned}$$

that is, the triangular lattices $T^t(n, m)$, $T^c(n, m)$, and $T^f(n, m)$ with toroidal, cylindrical, and free boundary conditions have the same asymptotic energy ($\approx 2.065mn$).

Proof. By definitions of $T^t(n, m)$, $T^c(n, m)$, and $T^f(n, m)$, $T^f(n, m)$ and $T^c(n, m)$ are spanning subgraph of $T^t(n, m)$. Moreover, almost all vertices of $T^f(n, m)$ or $T^c(n, m)$ are

6. Hence, by Corollary 2.5,

$$\lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(T^t(n, m))}{nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(T^c(n, m))}{nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(T^f(n, m))}{nm}.$$

It suffices to prove that

$$\lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(T^t(n, m))}{nm} = \frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{2\pi} |\cos x + \cos y + \cos(x+y)| dx dy \approx 2.065.$$

Let $A(C_m)$ be the adjacency matrix of the cycle C_m . By a suitable labelling of vertices of $T^t(n, m)$, the adjacency matrix $A(T^t(n, m))$ of $T^t(n, m)$ has the following form:

$$\begin{aligned} A(T^t(n, m)) &= \begin{pmatrix} A(C_m) & I_m + B_m & 0 & \cdots & 0 & I_m + B_m^T \\ I_m + B_m^T & A(C_m) & I_m + B_m & \cdots & 0 & 0 \\ 0 & I_m + B_m^T & A(C_m) & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A(C_m) & I_m + B_m \\ I_m + B_m & 0 & 0 & \cdots & I_m + B_m^T & A(C_m) \end{pmatrix}_{n \times n} \\ &= I_n \otimes A(C_m) + B_n \otimes (I_m + B_m) + B_n^T \otimes (I_m + B_m^T), \end{aligned}$$

where I_n is the identity matrix of order n , $M \otimes N$ denotes the tensor product of two matrices M and N , and

$$B_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \end{pmatrix}_{n \times n}.$$

Note that $A(C_n) = B_n + B_n^T$. Hence

$$A(T^t(n, m)) = I_n \otimes (B_m + B_m^T) + B_n \otimes (I_m + B_m) + B_n^T \otimes (I_m + B_m^T).$$

Let $\{1 = g^0, g^1, \dots, g^{n-1}\}$ be the cyclic group of order n . Obviously, $\rho : g^i \rightarrow B_n^i$ for $0 \leq i \leq n-1$ is a representation of this group. Note that the cyclic group of order n has exactly n (linear) characters χ_i ($i = 0, 1, \dots, n-1$) such that $\chi_i(g) = \omega_n^i$, where ω_n is the n th root of unitary. Hence there exists an invertible matrix $Q_n = (\frac{\omega_n^{ij}}{\sqrt{n}})_{0 \leq i, j \leq n-1}$ such that $Q_n^{-1} B_n Q_n = \text{diag}(1, \omega_n, \dots, \omega_n^{n-1}) =: D_n$. Since $B_n^T = B_n^{-1}$ and $Q_n^T = Q_n^{-1}$, $Q_n^{-1} B_n^T Q_n = \text{diag}(1, \omega_n^{-1}, \dots, \omega_n^{-(n-1)}) =: D_n^{-1}$. Hence

$$(Q_n^{-1} \otimes Q_m^{-1}) A(T^t(n, m)) (Q_n \otimes Q_m)$$

$$\begin{aligned}
&= (Q_n^{-1} \otimes Q_m^{-1})[I_n \otimes (B_m + B_m^T) + B_n \otimes (I_m + B_m) + B_n^T \otimes (I_m + B_m^T)](Q_n \otimes Q_m) \\
&= I_n \otimes (D_m + D_m^{-1}) + D_n \otimes (I_m + D_m) + D_n^{-1} \otimes (I_m + D_m^{-1}).
\end{aligned}$$

It is not difficult to see that $I_n \otimes (D_m + D_m^{-1}) + D_n \otimes (I_m + D_m) + D_n^{-1} \otimes (I_m + D_m^{-1})$ is a diagonal matrix whose diagonal entries are $\omega_m^j + \omega_m^{-j} + \omega_n^i + \omega_n^{-i} + \omega_n^i \omega_m^j + \omega_n^{-i} \omega_m^{-j} = 2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos(\frac{2i\pi}{n} + \frac{2j\pi}{m})$, $0 \leq i \leq n-1, 0 \leq j \leq m-1$. This implies that the eigenvalues of $A(T^t(n, m))$ are $2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos(\frac{2i\pi}{n} + \frac{2j\pi}{m})$, $0 \leq i \leq n-1, 0 \leq j \leq m-1$. By the definition of the energy,

$$\mathcal{E}(T^t(n, m)) = \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} |2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos(\frac{2i\pi}{n} + \frac{2j\pi}{m})|.$$

So

$$\begin{aligned}
\lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(T^t(n, m))}{nm} &= \lim_{n, m \rightarrow \infty} \frac{1}{nm} \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} |2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos(\frac{2i\pi}{n} + \frac{2j\pi}{m})| \\
&= \frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{2\pi} |\cos x + \cos y + \cos(x+y)| dx dy \approx 2.065
\end{aligned}$$

and we complete the proof of the theorem. $\square$




FIGURE 2. (a) The $3^3.4^2$ lattice $S^t(n, 2m)$ with toroidal boundary condition; (b) the $3^3.4^2$ lattice $S^c(n, 2m)$ with cylindrical boundary condition; (c) the $3^3.4^2$ lattice $S^f(n, 2m)$ with cylindrical boundary condition.

2.3. The $3^3.4^2$ lattice. The $3^3.4^2$ lattice $S^t(n, 2m)$ with toroidal boundary condition can be constructed by starting with a $2m \times n$ square lattice and adding a diagonal edge connecting the vertices in, say, the upper left to the lower right corners of each square in every other row as shown in Figure 2(a), where $a_1 = b_1, a_{2m} = b_1^*, a_1^* = b_n, a_{2m}^* = b_n^*$, and $(a_1, a_1^*), (a_2, a_2^*), \dots, (a_{2m}, a_{2m}^*), (b_1, b_1^*), (b_2, b_2^*), \dots, (b_n, b_n^*), (a_1, a_2^*), (a_3, a_4^*), \dots, (a_{2m-1}, a_{2m}^*)$ are edges in $S^t(n, 2m)$. If we delete edges $(b_1, b_1^*), (b_2, b_2^*), \dots, (b_n, b_n^*)$ from $S^t(n, 2m)$, the $3^3.4^2$ lattice $S^c(n, 2m)$ with cylindrical boundary condition is obtained (see Figure 2(b)).

If we delete edges $(a_1, a_1^*), (a_2, a_2^*), \dots, (a_{2m}, a_{2m}^*), (a_1, a_2^*), (a_3, a_4^*), \dots, (a_{2m-1}, a_{2m}^*)$ from $S^c(n, 2m)$, the $3^3.4^2$ lattice $S^f(n, 2m)$ with free boundary condition is obtained (see Figure 2(c)). The asymptotic number of spanning trees of $S^t(n, 2m)$ (resp. $S^c(n, 2m)$ and $T^f(n, 2m)$) was obtained by Chang and Wang [2] (resp. by Yan and Zhang [17]).

Theorem 2.9. *For the $3^3.4^2$ lattices $S^t(n, 2m)$, $S^c(n, 2m)$, and $S^f(n, 2m)$ with toroidal, cylindrical, and free boundary conditions,*

$$\begin{aligned} \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^t(n, 2m))}{2nm} &= \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^c(n, 2m))}{2nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^f(n, 2m))}{2nm} \\ &= \frac{1}{8\pi^2} \int_0^{2\pi} \int_0^{2\pi} (|2 \cos x + \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}| + \\ &\quad |2 \cos x - \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}|) dx dy \approx 1.8471, \end{aligned}$$

that is, the $3^3.4^2$ lattices $S^t(n, 2m)$, $S^c(n, 2m)$, and $S^f(n, 2m)$ with toroidal, cylindrical, and free boundary conditions have the same asymptotic energy ($\approx 3.6942mn$).

Proof. By definitions of $S^t(n, 2m)$, $S^c(n, 2m)$, and $S^f(n, 2m)$, $S^f(n, 2m)$ and $S^c(n, 2m)$ are spanning subgraphs of $S^t(n, 2m)$. Moreover, almost all vertices of $S^f(n, m)$ or $S^c(n, m)$ are of degree 5. Hence, by Corollary 2.5,

$$\lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^t(n, m))}{2nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^c(n, m))}{2nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^f(n, 2m))}{2nm}.$$

It suffices to prove that

$$\begin{aligned} \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^t(n, 2m))}{2nm} &= \frac{1}{8\pi^2} \int_0^{2\pi} \int_0^{2\pi} (|2 \cos x + \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}| + \\ &\quad |2 \cos x - \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}|) dx dy \approx 1.8471. \end{aligned}$$

Let $A(C_{2m})$ be the adjacency matrix of the cycle C_{2m} . By a suitable labelling of vertices of $S^t(n, 2m)$, the adjacency matrix $A(S^t(n, 2m))$ of $S^t(n, 2m)$ has the following form:

$$\begin{aligned} A(S^t(n, m)) &= \begin{pmatrix} A(C_{2m}) & I_{2m} + F_{2m} & 0 & \cdots & 0 & I_{2m} + F_{2m}^T \\ I_{2m} + F_{2m}^T & A(C_{2m}) & I_{2m} + F_{2m} & \cdots & 0 & 0 \\ 0 & I_{2m} + F_{2m}^T & A(C_{2m}) & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A(C_{2m}) & I_{2m} + F_{2m} \\ I_{2m} + F_{2m} & 0 & 0 & \cdots & I_{2m} + F_{2m}^T & A(C_{2m}) \end{pmatrix}_{n \times n} \\ &= I_n \otimes A(C_{2m}) + B_n \otimes (I_{2m} + F_{2m}) + B_n^T \otimes (I_{2m} + F_{2m}^T), \end{aligned}$$

where I_n is the identity matrix of order n , $M \otimes N$ denotes the tensor product of two matrices M and N , $B_n = (b_{ij})_{n \times n}$ such that $b_{ij} = 1$ if $(i, j) = (1, 2), (2, 3), \dots, (n-1, n), (n, 1)$ and $b_{ij} = 0$ otherwise, and $F_{2m} = (f_{ij})_{2m \times 2m}$ such that $f_{ij} = 1$ if $(i, j) = (2, 1), (4, 3), (6, 5), \dots, (2m, 2m-1)$ and $f_{ij} = 0$ otherwise.

Note that $A(C_{2m}) = I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + B_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + B_m^T \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Hence

$$\begin{aligned} A(S^t(n, 2m)) &= I_n \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + I_n \otimes B_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + I_n \otimes B_m^T \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ &\quad + B_n \otimes I_m \otimes \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} + B_n^T \otimes I_m \otimes \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

Let $Q_n = (\frac{\omega_n^{ij}}{\sqrt{n}})_{0 \leq i, j \leq n-1}$. Then $Q_n^{-1} B_n Q_n = \text{diag}(1, \omega_n, \dots, \omega_n^{n-1}) =: D_n$. Since $B_n^T = B_n^{-1}$ and $Q_n^T = Q_n^{-1}$, $Q_n^{-1} B_n^T Q_n = \text{diag}(1, \omega_n^{-1}, \dots, \omega_n^{-(n-1)}) =: D_n^{-1}$. Hence

$$\begin{aligned} &(Q_n^{-1} \otimes Q_m^{-1} \otimes I_2) A(S^t(n, 2m)) (Q_n \otimes Q_m \otimes I_2) \\ &= I_n \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + I_n \otimes D_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + I_n \otimes D_m^{-1} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ &\quad + D_n \otimes I_m \otimes \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} + D_n^{-1} \otimes I_m \otimes \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

It is not difficult to see that the above matrix is a block diagonal matrix whose diagonal blocks are $\begin{pmatrix} \omega_n^i + \omega_n^{-i} & 1 + \omega_n^{-i} + \omega_m^j \\ 1 + \omega_n^i + \omega_m^{-j} & \omega_n^i + \omega_n^{-i} \end{pmatrix}$, $0 \leq i \leq n-1, 0 \leq j \leq m-1$. Hence the eigenvalues of $A(S^t(n, 2m))$ are:

$$2 \cos \frac{2i\pi}{n} \pm \sqrt{3 + 2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos \left(\frac{2i\pi}{n} + \frac{2j\pi}{m} \right)}, \quad 0 \leq i \leq n-1, 0 \leq j \leq m-1.$$

Set

$$\begin{aligned} \lambda_1(i, j) &= 2 \cos \frac{2i\pi}{n} + \sqrt{3 + 2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos \left(\frac{2i\pi}{n} + \frac{2j\pi}{m} \right)}, \\ \lambda_2(i, j) &= 2 \cos \frac{2i\pi}{n} - \sqrt{3 + 2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos \left(\frac{2i\pi}{n} + \frac{2j\pi}{m} \right)}. \end{aligned}$$

By the definition of the energy,

$$\mathcal{E}(S^t(n, 2m)) = \sum_{i=0}^{n-1} \sum_{j=0}^{m-1} (|\lambda_1(i, j)| + |\lambda_2(i, j)|).$$

So

$$\lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(S^t(n, 2m))}{2nm} = \frac{1}{8\pi^2} \int_0^{2\pi} \int_0^{2\pi} (|2 \cos x + \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}| + |2 \cos x - \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)}|) dx dy \approx 1.8471$$

and we complete the proof of the theorem. $\square$




FIGURE 3. (a) The hexagonal lattice $H^t(n, m)$ with toroidal boundary condition; (b) the hexagonal lattice $H^c(n, m)$ with cylindrical boundary condition; (c) the hexagonal lattice $H^f(n, m)$ with cylindrical boundary condition.

2.4. The hexagonal lattice. The hexagonal lattices with toroidal and cylindrical boundary conditions, denoted by $H^t(n, m)$ and $H^c(n, m)$, are illustrated in Figure 3(a) and Figure 3(b), where $(a_1, b_1), (a_2, b_2), \dots, (a_{m+1}, b_{m+1}), (a_1, c_1^*), (c_1, c_2^*), (c_2, c_3^*), \dots, (c_{n-1}, c_n^*), (c_n, b_{m+1})$ are edges in $H^t(n, m)$, and $(a_1, b_1), (a_2, b_2), \dots, (a_{m+1}, b_{m+1})$ are edges in $H^c(n, m)$. If we delete edges $(a_1, b_1), (a_2, b_2), \dots, (a_{m+1}, b_{m+1})$ from $H^c(n, m)$, then the hexagonal lattice, denoted by $H^f(n, m)$, with free boundary condition is obtained (see Figure 3(c)). The asymptotic number of perfect matchings of $H^t(n, m)$ can be found in Wu [15]. The asymptotic number of spanning trees of $H^t(n, m)$ (resp. $H^c(n, m)$ and $H^f(n, m)$) was obtained by Shrock and Wu [14] (resp. by Yan and Zhang [17]).

Theorem 2.10. *For the hexagonal lattices $H^t(n, m)$, $H^c(n, m)$, and $H^f(n, m)$ with toroidal, cylindrical, and free boundary conditions,*

$$\begin{aligned} \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(H^t(n, m))}{2nm} &= \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(H^c(n, m))}{2nm} = \lim_{n,m \rightarrow \infty} \frac{\mathcal{E}(H^f(n, m))}{2nm} \\ &= \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)} dx dy \approx 1.5746, \end{aligned}$$

that is, the hexagonal lattices $H^t(n, m)$, $H^c(n, m)$, and $H^f(n, m)$ with toroidal, cylindrical, and free boundary conditions have the same asymptotic energy ($\approx 3.1492mn$).

Proof. By definitions of $H^t(n, m)$, $H^c(n, m)$, and $H^f(n, m)$, $H^f(n, m)$ and $H^c(n, m)$ are spanning subgraphs of $H^t(n, m)$. Moreover, almost all vertices of $H^f(n, m)$ or $H^c(n, m)$ are 3. Hence, by Corollary 2.5,

$$\lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(H^t(n, m))}{2nm} = \lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(H^c(n, m))}{2nm} = \lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(H^f(n, m))}{2nm}.$$

It suffices to prove that

$$\lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(H^t(n, m))}{2nm} = \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} \sqrt{3 + 2\cos x + 2\cos y + 2\cos(x+y)} dx dy \approx 1.5746.$$

Let $A(C_{2m})$ be the adjacency matrix of the cycle C_{2m} . By a suitable labelling of vertices of $H^t(n-1, m-1)$, the adjacency matrix $A(H^t(n-1, m-1))$ of $H^t(n-1, m-1)$ has the following form:

$$\begin{aligned} A(H^t(n-1, m-1)) &= \begin{pmatrix} A(C_{2m}) & F_{2m} & 0 & \cdots & 0 & F_{2m}^T \\ F_{2m}^T & A(C_{2m}) & F_{2m} & \cdots & 0 & 0 \\ 0 & F_{2m}^T & A(C_{2m}) & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A(C_{2m}) & F_{2m} \\ F_{2m} & 0 & 0 & \cdots & F_{2m}^T & A(C_{2m}) \end{pmatrix}_{n \times n} \\ &= I_n \otimes A(C_{2m}) + B_n \otimes F_{2m} + B_n^T \otimes F_{2m}^T, \end{aligned}$$

where I_n is the identity matrix of order n , $M \otimes N$ denotes the tensor product of two matrices M and N , $B_n = (b_{ij})_{n \times n}$ such that $b_{ij} = 1$ if $(i, j) = (1, 2), (2, 3), \dots, (n-1, n), (n, 1)$ and $b_{ij} = 0$ otherwise, and $F_{2m} = (f_{ij})_{2m \times 2m}$ such that $f_{ij} = 1$ if $(i, j) = (2, 1), (4, 3), (6, 5), \dots, (2m, 2m-1)$ and $f_{ij} = 0$ otherwise.

Note that $A(C_{2m}) = I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + B_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + B_m^T \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Hence

$$\begin{aligned} A(H^t(n-1, m-1)) &= I_n \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + I_n \otimes B_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + I_n \otimes B_m^T \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ &\quad + B_n \otimes I_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + B_n^T \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Let $Q_n = (\frac{\omega_n^{ij}}{\sqrt{n}})_{0 \leq i, j \leq n-1}$. Then $Q_n^{-1} B_n Q_n = \text{diag}(1, \omega_n, \dots, \omega_n^{n-1}) =: D_n$. Since $B_n^T = B_n^{-1}$ and $Q_n^T = Q_n^{-1}$, $Q_n^{-1} B_n^T Q_n = \text{diag}(1, \omega_n^{-1}, \dots, \omega_n^{-(n-1)}) =: D_n^{-1}$. Hence

$$(Q_n^{-1} \otimes Q_m^{-1} \otimes I_2) A(H^t(n-1, m-1)) (Q_n \otimes Q_m \otimes I_2)$$

$$\begin{aligned}
&= I_n \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + I_n \otimes D_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + I_n \otimes D_m^{-1} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
&\quad + D_n \otimes I_m \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + D_n^{-1} \otimes I_m \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.
\end{aligned}$$

It is not difficult to see that the above matrix is a block diagonal matrix whose diagonal blocks are $\begin{pmatrix} 0 & 1 + \omega_n^{-i} + \omega_m^j \\ 1 + \omega_n^i + \omega_m^{-j} & 0 \end{pmatrix}$, $0 \leq i \leq n-1, 0 \leq j \leq m-1$. Hence the eigenvalues of $A(H^t(n-1, m-1))$ are: $\pm \sqrt{3 + 2 \cos \frac{2i\pi}{n} + 2 \cos \frac{2j\pi}{m} + 2 \cos \left(\frac{2i\pi}{n} + \frac{2j\pi}{m} \right)}$, $0 \leq i \leq n-1, 0 \leq j \leq m-1$.

By the definition of the energy,

$$\mathcal{E}(H^t(n, m)) = 2 \sum_{i=0}^n \sum_{j=0}^m \sqrt{3 + 2 \cos \frac{2i\pi}{n+1} + 2 \cos \frac{2j\pi}{m+1} + 2 \cos \left(\frac{2i\pi}{n+1} + \frac{2j\pi}{m+1} \right)}.$$

So

$$\lim_{n, m \rightarrow \infty} \frac{\mathcal{E}(H^t(n, m))}{2(n+1)(m+1)} = \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} \sqrt{3 + 2 \cos x + 2 \cos y + 2 \cos(x+y)} dx dy \approx 1.5746$$

and we complete the proof of the theorem. $\square$

We would like to point out that the result of hexagonal lattice with toroidal boundary condition has been obtained in [8] by a different approach.

Remark 2.11. For the triangular, $3^3.4^2$, and hexagonal lattices, we have considered the three boundary conditions: the toroidal, cylindrical, and free boundary conditions in Theorem 2.8-2.10, respectively. By a similar idea, we can consider another two boundary conditions: the Mobius-band and Klein-bottle boundary conditions, and show that the triangular (resp. $3^3.4^2$ and hexagonal) lattices with these five boundary conditions have the same asymptotic energy.

3. CONCLUDING REMARKS

In this paper, we showed that for many types of lattices the energy per vertex of the plane lattices is independent of the boundary conditions. It is no difficulty to see that the conclusion is true in general. In fact our approach can be used widely. By using this conclusion we can convert some harder problem to easy one and get some results simultaneously. For example dealing with the problem of the asymptotic energy of the hexagonal lattice with the free boundary is not an easy task but we deduced it in a simple way. On the other hand, for the entropy of dimers the result is not true. In fact, Yan, Yeh,

and Zhang [16] showed that for the dimer problem the hexagonal lattices with cylindrical and toroidal boundary have different entropies.

References

- [1] N. J. Calkin and H. S. Wilf, *The number of independent sets in a grid graph*, SIAM J. DISCRETE MATH. Vol. 11 (1998) 54C-60.
- [2] S.-C. Chang and W. Wang, *Spanning trees on lattices and integral identities*, J. Phys. A: Math. Gen., 39 (2006) 10263–10275.
- [3] J. Day, W. So, *Singular value inequality and graph energy change*, Electronic Journal of Linear Algebra, 16 (2007), 291–299.
- [4] J. Day, W. So, *Graph energy change due to edge deletion*, Linear Algebra and its Applications, 428 (2008), 2070–2078.
- [5] M. E. Fisher, *Statistical mechanics of dimers on a plane lattice*, Phys. Rev., 124(1961), 1664–1672.
- [6] I. Gutman, *The energy of a graph*, Ber. Math. -Statist. Sect. Forschungszentrum Graz, 103 (1978), 1–22.
- [7] I. Gutman, *The energy of a graph: old and new results*, Algebraic Combinatorics and Applications (Gossweinstein, 1999), Springer, Berlin, 2001, pp. 196–211.
- [8] J. H. Hua, *The spectra of some lattice graphs on surfaces*, Dissertation of Lanzhou University, 2008.
- [9] P. W. Kasteleyn, *The statistics of dimers on a lattice I: The number of dimer arrangements on a quadratic lattice*, Physica, 27(1961), 1209–1225.
- [10] P. W. Kasteleyn, *Dimer statistics and phase transitions*, J. Math. Phys., 4(1963), 287–293.
- [11] P. W. Kasteleyn, *Graph Theory and Crystal Physics*, Graph Theory and Theoretical Physics (F. Harary, ed.), Academic Press, 1967, 43–110.
- [12] J. H. Koolen, V. Moulton, *Maximal energy graphs*, Adv. Appl. Math. 26 (2001), 47–52.
- [13] H. N. V. Temperley, M. E. Fisher, *Dimer problem in statistical mechanics—an exact result*, Philosophical Magazine, 6(1961), 1061–1063.
- [14] R. Shrock, F. Y. Wu, *Spanning trees on graphs and lattices in d dimensions*, J. Phys. A: Math. Gen., 33 (2000), 3881–3902.

- [15] F. Y. Wu, *Dimers on two-dimensional lattices*, Intern. J. Modern Phys. B, 32 (2006), 5357–5371.
- [16] W. G. Yan, Y.-N. Yeh, and F. J. Zhang, *Dimer problem on the cylinder and torus*, Physica A: Statistical Mechanics and its Applications, 387 (2008), 6069–6078.
- [17] W. G. Yan, F. J. Zhang, *Enumeration of spanning trees of some symmetric graphs*, Preprint.

SCHOOL OF SCIENCES, JIMEI UNIVERSITY, XIAMEN 361021, CHINA

E-mail address: weigenyan@263.net

CORRESPONDING AUTHOR, DEPARTMENT OF MATHEMATICS, WEST VIRGINIA UNIVERSITY, MORGANTOWN, WV 26506-6310, USA

E-mail address: zhzhang@math.wvu.edu